SPACEFLIGHT NOW The leading source for online space news breaking news

HOME = CURRENT MISSION = PLUS = NEWS ARCHIVE = LAUNCH SCHEDULE = MISSION REPORT = ASTRONOMY NOW = STORE

Supermassive black hole at core of skinny galaxy

UNIVERSITY OF CALIFORNIA-BERKELEY NEWS RELEASE Posted: April 12, 2003

Scientists have uncovered a supermassive black hole at the core of a svelte, spiral galaxy, a finding that questions a recently devised rule of thumb in which only galaxies with bulging cores have such black holes.

Dr. Alex Filippenko, professor of astronomy at the University of California, Berkeley, and Dr. Luis Ho, an astronomer at the Observatories of the Carnegie Institution of Washington, in Pasadena, discuss these results in the May 1, 2003, issue of The Astrophysical Journal Letters.

The scientists determined that galaxy NGC 4395, a flat "pure-disk" galaxy with no central bulge, has a central black hole approximately 10,000 to 100,000 times the mass of our sun. This suggests that other pure-disk galaxies, thought to be devoid of supermassive black holes, may indeed have one lurking within quite possibly the featherweights of the supermassive black hole club.

"The supermassive black hole in NGC 4395 is the smallest one yet found in the center of a galaxy," said Filippenko. "This would be consistent with the galaxy having a small bulge. However, the bulge is not just small, it seems to be nonexistent."

Supermassive black holes typically range from millions to billions of times the mass of the Sun, dwarfing the more common stellar black holes that are created by the runaway gravitational collapse of the cores of massive stars.

NGC 4395 is 11 million light years away in the Northern Hemisphere constellation of Canes Venatici ("the Hunting Dogs"). The galaxy has long been known to emit significant amounts of light, including visible light and X rays, from its central core region. This is a telltale sign of the presence of a central, giant black hole actively sucking in enough matter to create a swirling, superheated accretion disk. Yet no supermassive black hole had been detected.

Filippenko and Ho used the Keck I 10-meter telescope on Mauna Kea in Hawaii and the Japanese-U.S. ASCA X ray telescope to determine that NGC 4395 indeed has a supermassive black hole, albeit an unusually light one.

Starry Night

The highly acclaimed Starry Night range of software is now available from the Astronomy Now Store. Put the universe on your desktop.



■ <u>U.S. STORE</u>

2002 in review

NIEWA/

Astronomy Now and Spaceflight Now present 2002: a Year in Space -- a stunning month-by-



month pictorial record of the space events and discoveries during 2002. It is a musthave for all space enthusiasts!

- <u>U.S. STORE</u>
- U.K. & WORLDWIDE STORE

The ultimate Apollo 11 DVD

NEW 3-DISC EDITION This exceptional chronicle of the historic Apollo 11 lunar landing mission features



new digital transfers of film and television coverage unmatched by any other.

- U.S. STORE
- U.K. & WORLDWIDE STORE

Get e-mail updates

Sign up for our NewsAlert service and have the latest news in astronomy and space e-mailed direct to your desktop (privacy note: your e-mail address will not be used for any other purpose).

Enter your e-mail address:

Subscribe

"Here is an example of a massive black hole that is low in comparison to all previously reported supermassive ones, but it is definitely much more massive than stellar-class black holes, and is located in a galaxy that has no bulge," Ho said. "Thus, having a well-developed bulge is evidently not a necessary condition for the formation of massive, central black holes."

Filippenko and Ho confirmed the mass range using the velocity dispersion technique for estimating the mass of black holes, and in the process found that this powerful tool for inferring black hole mass works on scales far lower than previously thought.

Velocity dispersion refers to how fast, on average, the stars in a galactic core region are orbiting about the central black hole. It is akin to taking an average speed of a swarm of bees circling about their hive on paths (orbits) of differing size and orientation. Although predicted theoretically back in the 1990s, confirmation of a very tight correlation between velocity dispersion and supermassive black hole mass came in 2000. But until now, that correlation was always assumed to apply to black holes weighing in at millions or billions of solar masses. Using this technique on the NGC 4395, Filippenko and Ho found its mass to be 66,000 solar masses - nearly in the middle of the range found using X-ray luminosity.

This means that the black hole mass/velocity dispersion relation may apply even on scales of relatively small star clusters that contain a central, relatively light, supermassive black hole, Ho said.

The scientists also suggest that NGC 4395 may represent a unique step in the evolution of supermassive black holes, in which a bulge will develop as the black hole grows. The velocity dispersion measurements were based on a star cluster near the central black hole, a possible indication that a bulge may someday form.

Hubble Calendar

NEW! This remarkable calendar features stunning images of



planets, stars, gaseous nebulae, and galaxies captured by NASA's orbiting Hubble Space Telescope.

- <u>U.S. STORE</u>
- U.K. & WORLDWIDE STORE

Hubble Posters

Stunning posters featuring images from the Hubble Space Telescope and worldrenowned



astrophotographer David Malin are now available from the Astronomy Now Store.

- U.S. STORE
- U.K. & WORLDWIDE STORE

Baseball caps

The NASA
"Meatball" logo
appears on a
series of stylish
baseball caps
available now
from the Astronomy Now
Store.

- U.S. STORE
- U.K. & WORLDWIDE STORE

Space station sweatshirt

NEW!Our cozy
Space Station
cadet collar
pullover
sweatshirt is
perfect for
keeping warm on a cold
winter night.

- U.S. STORE
- WORLDWIDE STORE

 $\frac{INDEX \mid PLUS \mid BREAKING \ NEWS \mid LAUNCH \ SCHEDULE}{MISSION \ REPORT \mid FEATURES \mid ASTRONOMY \ NOW \mid STORE}$ ADVERTISE

© 2003 Pole Star Publications Ltd